

OHSTPY-HEP-T-01-025

August 2001

Yukawa Coupling Unification in SO(10) Supersymmetric Grand Unified Theories^{abc}

Radovan Dermíšek

*Department of Physics, The Ohio State University,
174 W. 18th Ave., Columbus, Ohio 43210*

Abstract

We report on a preferred region of supersymmetric parameter space by t , b , τ Yukawa unification. We find a narrow region survives for $\mu > 0$ (suggested by $b \rightarrow s\gamma$ and the anomalous magnetic moment of the muon) with $A_0 \sim -2 m_{16}$, $m_{16} > 1200$ GeV, $\mu, M_{1/2} \ll m_{16}$ and $\tan \beta \sim 50 \pm 2$. Demanding Yukawa unification thus makes definite predictions for Higgs and sparticle masses.

^aTalk given at the 9th International Conference on Supersymmetry and Unification of Fundamental Interactions SUSY'01, June 11–17, Dubna, Russia.

^bThis talk is based on the work done in collaboration with T. Blažek and S. Raby.

^cThis work was supported in part by DOE grant DOE/ER/01545-820.

YUKAWA COUPLING UNIFICATION IN $SO(10)$ SUPERSYMMETRIC GRAND UNIFIED THEORIES

RADOVAN DERMIŠEK

*Department of Physics, The Ohio State University,
174 W. 18th Ave., Columbus, OH 43210, USA
E-mail: dermisek@pacific.mps.ohio-state.edu*

We report on a preferred region of supersymmetric parameter space by t , b , τ Yukawa unification. We find a narrow region survives for $\mu > 0$ (suggested by $b \rightarrow s\gamma$ and the anomalous magnetic moment of the muon) with $A_0 \sim -2m_{16}$, $m_{16} > 1200$ GeV, $\mu, M_{1/2} \ll m_{16}$ and $\tan\beta \sim 50 \pm 2$. Demanding Yukawa unification thus makes definite predictions for Higgs and sparticle masses.

The nicest features of supersymmetric [SUSY] grand unified theories [GUTs] are the explanation of charge assignments of quarks and leptons and the prediction of the weak mixing angle from the unification of gauge couplings at the GUT scale $M_G \sim 3 \times 10^{16}$ GeV. Furthermore, there is also a very interesting possibility that the Yukawa couplings of the third generation particles unify at the same scale. The unification of Yukawa couplings of t , b , τ is very well motivated in $SO(10)$ models where the third generation standard model [SM] fermions and the righthanded neutrino sit in one **16** dimensional spinor representation and (in the simplest version) the two Higgs doublets of minimal supersymmetric standard model are contained in one **10** dimensional fundamental representation. The Yukawa couplings are given by a single term in the superpotential $W = \lambda \mathbf{16} \mathbf{10} \mathbf{16}$ which leads to $\lambda_t = \lambda_b = \lambda_\tau = \lambda$ at the GUT scale. Besides λ and M_G there is one more parameter crucial for fermion masses – $\tan\beta$, the ratio of vevs of two Higgses. Since M_G is already fixed by gauge coupling unification there are just two parameters to play with and it is non-trivial that one can describe three masses by one Yukawa coupling. In what follows we specify the SUSY breaking scenario and comment on alternatives, we show that there is a preferred region of SUSY parameter space where Yukawa unification works, and give predictions for Higgs and SUSY spectra from this specific region. For a more detailed discussion see Ref.¹.

In our analysis we assume following boundary conditions for soft SUSY breaking terms at M_G : universal squark and slepton masses – m_{16} ; universal gaugino masses – $M_{1/2}$; non-universal Higgs H_u , H_d masses; and universal trilinear couplings – A . The special case $m_{16} = m_{H_u} = m_{H_d}$ corresponds to the MSUGRA scenario. However it is easier to obtain radiative electroweak symmetry breaking [REWSB] in large $\tan\beta$ regime when Higgs masses are

split ($m_{H_u} < m_{H_d}$). Another alternative would be to consider D-term contribution to scalar masses from breaking the rank of $SO(10)$: $m_{(H_u, H_d)}^2 = m_{10}^2 \mp 2D_X$, $m_Q^2 = m_E^2 = m_U^2 = m_{16}^2 + D_X$, $m_D^2 = m_L^2 = m_{16}^2 - 3D_X$. In this case REWSB is possible if $D_X > 0$ and the Yukawa unification can be achieved only for $\mu < 0$.²

The Yukawa unification is easy to satisfy with $\mu < 0$, because reasonable fits to b quark mass prefer negative SUSY threshold corrections. In most region of SUSY parameter space these are dominated by gluino contribution which are negative for negative μ . However, the discrepancy between SM and experimental values of anomalous magnetic moment of muon can be explained by SUSY only if $\mu > 0$.³ Also $b \rightarrow s\gamma$ for large $\tan\beta$ strongly prefers $\mu > 0$.⁴

In this analysis we are looking for the region of SUSY parameter space with $\mu > 0$ consistent with Yukawa unification. We start with 11 input parameters at the GUT scale: M_G , α_G , ϵ_3 , (ϵ_3 is a correction to α_3 at the GUT scale) which are fixed by gauge coupling unification; m_{H_u} , m_{H_d} which are constrained by requiring correct REWSB; λ , $\tan\beta$ which are constrained by Yukawa unification; and remaining parameters μ , $M_{1/2}$, m_{16} , A affect Yukawa unification through RG running and weak scale threshold corrections. We scan through these parameters to find the region where Yukawa unification can be satisfied. The χ^2 function includes 9 observables: 6 precision electroweak data α_{EM} , G_μ , $\alpha_s(M_Z)$, M_Z , M_W , ρ_{NEW} and the 3 fermion masses M_{top} , $m_b(m_b)$, M_τ .

Fig. 1 shows contours of constant χ^2 (dashed lines) for $m_{16} = 2000$ GeV. These contours basically correspond to contours of constant m_b (not shown in the Fig. 1)⁵ which typically gives the largest contribution to χ^2 . The contour of $\chi^2 = 1$ roughly corresponds to $m_b(m_b) \sim 4.25$ GeV and $\chi^2 = 3$ to $m_b(m_b) \sim 4.4$ GeV. Fig. 1 (left) also shows contours of constant SUSY threshold corrections to m_b . We see that the acceptable fits require $\Delta m_b \leq -2\%$.

We find acceptable fits ($\chi^2 < 3$) for $m_{16} > 1200$ GeV, $A \sim -2m_{16}$, μ , $M_{1/2} \ll m_{16}$ and $\tan\beta \sim 50 \pm 2$. Why does Yukawa unification only work in this narrow region of SUSY parameter space? How is it possible to get negative threshold corrections with positive μ ? The dominant corrections to m_b comes from gluino and chargino diagrams. These are proportional to $\tan\beta$ and can naturally be as large as 40 %:

$$\Delta m_b^{\tilde{g}} \approx \frac{2\alpha_3}{3\pi} \frac{\mu m_{\tilde{g}}}{m_b^2} \tan\beta \quad \text{and} \quad \Delta m_b^{\tilde{\chi}^\pm} \approx \frac{\lambda_t^2}{16\pi^2} \frac{\mu A_t}{m_t^2} \tan\beta. \quad (1)$$

Gluino corrections are positive for $\mu > 0$ while chargino corrections are negative, since A_t runs to an infrared fixed point $\propto -M_{1/2}$. Hence in order to have a negative total correction to m_b , chargino corrections must dominate. This




Figure 1. Contours of constant Δm_b in % (left) and m_{h0} (GeV) (right) for $m_{16} = 2000$ GeV with the constant χ^2 contours overlayed (dashed lines). These correspond to $\chi^2 = 1, 2, 3$ and 5 in this order from the center of the picture out. The shaded region is excluded by chargino mass limit $m_{\tilde{\chi}^\pm} > 103$ GeV.

can be achieved if $A_t \ll 0$ and $m_{\tilde{t}} < m_{\tilde{b}}$. Large values of m_{16} are necessary to obtain large enough splitting between stop and sbottom. We typically find $m_{\tilde{b}_1} \sim 3 m_{\tilde{t}_1}$. This also explains why D-term splitting scenario with $D_X > 0$ is disfavoured. It makes the sbottom lighter then the stop already at the GUT scale and leads to $m_{\tilde{b}_1} \leq m_{\tilde{t}_1}$.

The typical spectrum for this region of SUSY parameter space is: quite light stop $\tilde{t}_1 \sim 150 - 250$ GeV, chargino $\tilde{\chi}^\pm \sim 100 - 250$ GeV and scalars of the first two generations above 1 TeV. Finally in Fig. 1 (right) we see that Yukawa unification prefers light Higgs mass in a narrow range, 110 - 118 GeV.

The range of SUSY parameters with $m_{16} > 1200$ GeV and $M_{1/2} \ll m_{16}$ is also preferred by nucleon decay experiments.⁶ However, large values of $m_{16} \geq 1200$ GeV lead to very small values of $a_\mu^{NEW} \leq 16 \times 10^{-10}$.

References

1. T. Blažek, R. Dermíšek and S. Raby, hep-ph/0107097.
2. H. Baer *et al*, *Phys. Rev. D* **61**, 111701 (2000).
3. See for example, R. Arnowitt, this volume.
4. See for example, W. de Boer, this volume.
5. For more figures see transparencies of this talk at www.susy.dubna.ru; see also S. Raby, this volume.
6. R. Dermíšek, A. Mafi and S. Raby, *Phys. Rev. D* **63**, 035001 (2001).